

# **KANGAROO SURVEY AND HABITAT ASSESSMENT**

**Scholars Drive – Link Road, Sippy Downs**

**For**

**Sunshine Coast Council**



**Native Foresters**

**Natural Resource Management Consultants**

Email: [simon@nativeforesters.com](mailto:simon@nativeforesters.com)

Mob: 0407 757 872

Document No: 221021-KHA

Date: 22<sup>nd</sup> October 2021

**DISCLAIMER**

The authors do not accept any form of liability, be it contractual, tortious, or otherwise, for the contents of this document or for any consequences arising from its use or any reliance placed upon it. This report has been compiled on the basis of existing literature, information, and a targeted survey. Native Foresters does not undertake responsibility arising in any way to any persons in respect of the data, errors, or omissions arising, through misinterpretation of information, negligence or otherwise, however caused. Native Foresters is an independent consulting company undertaking projects in Natural Resource Management.

All work conducted as part of this survey was conducted under the DES Scientific Purposes Permit number WISP 10189211 and DEEDI Animal Ethics Committee number CA 2020/08/1398 and in accordance with the relevant regulations.

**STUDY TEAM**

Simon McVerry (BASc – NRM) - Project Director  
Scott Burnett (PhD) – Fieldwork and reporting

**DOCUMENT CONTROL**

<b>Project:</b>	Scholars Drive Link Road
<b>Document:</b>	Kangaroo Survey and Habitat Assessment
<b>Document No.:</b>	221021-KHA
<b>Client:</b>	Sunshine Coast Council
<b>Project Manager:</b>	Yasmin Beavis

## Table of Contents:

1.0	INTRODUCTION .....	4
1.1	Scope.....	4
1.2	Objectives .....	4
1.3	Relevant Legislation and Guidelines .....	4
1.4	Proposed Works .....	7
1.5	Site Description .....	7
2.0	HABITAT ASSESSMENT .....	8
2.1	Methodology .....	8
2.2	Results.....	8
2.2.1	Desktop.....	8
2.2.2	Site Survey .....	9
2.2.3	Habitat Assessment .....	9
2.3	Kangaroo Research at USC.....	10
3.0	POTENTIAL IMPACTS OF WORKS .....	11
3.1	Kangaroo population .....	11
3.2	Sustainable Population Limit for USC.....	11
4.0	KANGAROO MANAGEMENT PLAN .....	12
4.1	Design Considerations .....	12
4.2	Construction Considerations .....	12
4.3	Post-Construction Considerations .....	13
5.0	SUMMARY .....	14
	REFERENCES.....	15
	APPENDIX A – INCIDENT AND COMPLAINT FORM.....	16
	APPENDIX B – Fauna recorded on USC campus .....	17
	APPENDIX C - RELOCATION PROCEDURE.....	22

## Figures:

Figure 1:	Link Road alignment and survey area.....	5
Figure 2:	Link Road location within USC Campus.....	5
Figure 3:	Proposed Works Alignment.....	6
Figure 4:	Scholars Drive link looking east towards the Parkville Road connection.....	7
Figure 5:	Scholars Drive link looking north towards the Scholars Drive connection.....	9

## 1.0 INTRODUCTION

Sunshine Coast Council have engaged Native Foresters to undertake an assessment of an Eastern Grey Kangaroo (*Macropus giganteus*) population within an area proposed for construction of a link road within the University of the Sunshine Coast (USC) Campus. The proposed link road is between Scholars Road and Parkville Street, Sippy Downs and its objective is to relieve traffic congestion related to adjacent schools.

### 1.1 Scope

This Kangaroo Survey and Habitat Assessment has been prepared for Sunshine Coast Council and relates to the proposed construction of a link road, connecting Scholars Drive to the northern end of Parkville Street, Sippy Downs. Refer to **Figure 1** for the survey area, **Figure 2** for the location of works on the university campus, **Figure 3** showing proposed works.

This report outlines the findings of the site survey and habitat assessment and proposes mitigation measures to ensure that impacts on Kangaroo populations and their habitats are avoided and/or minimised.

### 1.2 Objectives

The objectives of the FMP are as follows:

- To assess habitat value of works corridor of the proposed link road.
- To determine potential impacts to the Eastern Grey Kangaroo (*Macropus giganteus*) population in the vicinity as a result of the proposed works.
- To recommend mitigation and management actions that address and manage risks to the existing Kangaroo population within the subject site, including risks associated with vegetation clearing, earthworks, impacts to habitat areas and fauna rescue provisions.

### 1.3 Relevant Legislation and Guidelines

The following legislation and guidelines were considered relevant to the preparation of this FMP:

- *Nature Conservation Act 1992* (NCA) and Regulations – State
- *Animal Care and Protection Act 2001* - State
- *Code of Practice: Care of Sick, Injured or Orphaned Protected Animals in Queensland*.



**Figure 1: Link Road alignment and survey area**



**Figure 2: Link Road location within USC Campus**






Figure 3: Proposed Works Alignment

## 1.4 Proposed Works

The proposed Scholars Drive Link Road is an approximately 250m long and 3m wide concrete vehicle access road that will operate one way, at 20km/h during peak school drop off and pickup times only to relieve congestion. The road will open to traffic via an automatic boom gate during peak times in the morning and afternoon. It will function as a pedestrian pathway at all other times.

Seventeen, 1m high bollard lights will be installed at 15m intervals along the uphill side of the link road. It's proposed that the bollard lights will be active for approximately 9 hours each night and will have a level of backward shielding fitted to minimise light spill towards the adjacent residences.

## 1.5 Site Description

The works area is a 3000m<sup>2</sup> corridor of lawn and slashed introduced grasses on the south-western boundary of the University of the Sunshine Coast (USC) Sippy Downs campus. The link road route is sited entirely within an established strip of mown lawn/paddock which serves as a buffer around the inside of the USC boundary in this area.

This grassed buffer area is sandwiched between an extensive, regenerating wetland system on the USC campus site, and the back fences of houses within the suburb of Chancellor Park. The link road will be located within several metres of the wetland, and up to 10m from the boundary fence. A high timber fence marks the boundary of the USC property and the adjacent suburb as shown in **Figure 4**.



**Figure 4: Scholars Drive link looking east towards the Parkville Road connection**

## 2.0 HABITAT ASSESSMENT

### 2.1 Methodology

A site field survey was conducted on the 5<sup>th</sup> October 2021. Weather was as follows:

- Low humidity, with approximate temperature range of 12.5 – 27 degrees Celsius.
- The Caloundra racecourse weather station recorded 0 mm of rain on the 5<sup>th</sup> October and a total of 11.4 mm of rain fell in the week preceding the survey.

The following methods were used to determine the habitat values for native fauna:

- A desktop review was undertaken to assist in determining the site's ecological attributes. The Queensland Government's Regulated Vegetation Management Map (DR) was consulted to determine the mapped vegetation category and the presence of Essential Habitat. A search of the Queensland Government's (DES) WildNET database was conducted to generate a list of confirmed rare and threatened species records for the survey area. The School of Science, Technology and Engineering research results and papers on Eastern grey kangaroos on the USC campus were consulted.
- A visual and active search for signs of Eastern Grey Kangaroo use or habitation in the proposed link road corridor.
- A fauna habitat assessment based on the suitable breeding habitats present, the listed species known to occur or potentially occurring within the locality and the occurrence of specific habitat features appropriate for these species.

## 2.2 Results

### 2.2.1 Desktop

Following database and document reviews:

- The works corridor does not contain remnant vegetation on the RVMM.
- The Wildnet database lists Eastern Grey Kangaroo (*Macropus giganteus*) as being recorded within 1 km of the works corridor, as well as five EVNT listed fauna species within 1km of the site:
  - *Litoria freycineti* (Wallum rocketfrog) – *Vulnerable*
  - *Litoria olongburensis* (Wallum sedgefrog) – *Vulnerable*
  - *Hirundapus caudacutus* (White-throated needletail) – *Vulnerable*
  - *Phascolarctos cinereus* (Koala) – *Vulnerable*
  - *Pteropus poliocephalus* (Grey-headed flying-fox) – *Vulnerable under the EPBC Act*
- The works are within the Mooloolo River catchment and are just south of a mapped low-risk waterway under DAF mapping.
- The vegetation and habitats of the USC campus are well studied and support the breeding, foraging and habitation requirements of a diverse assemblage of almost 200 native fauna including 27 mammal species, 129 bird species, 20 reptile species, 8 amphibian species and 7 fish species (unpubl. data collected by S. Burnett, Jill Chamberlain, Gerard Mills 2007-2020). Refer to **Appendix B**.



## 2.2.2 Site Survey

A site survey was conducted in early October 2021. Around a dozen eastern grey kangaroo (*Macropus giganteus*) scats of different sizes were found throughout the study area, indicating that it is presently used as a foraging and/or movement pathway by a number of kangaroos.

The dry, elevated area of the proposed link road, mainly consisting of an unwatered lawn of couch and kikuyu grass (refer **Figure 4**). A short section of the route, at the Scholars Drive end, encompasses slashed *Setaria* tussocks which are a higher quality forage for the Eastern Grey Kangaroos than at other parts of the work's alignment as shown in **Figure 5**. The site survey coincided with an extended period of dry weather and the relative palatability of grasses along either section of the link may change under different times of the year. The alignment of the link road avoids any native vegetation. The current alignment will create a 0.13 ha island of lawn between the road and the USC/Chancellor Park boundary fence.



**Figure 5: Scholars Drive link looking north towards the Scholars Drive connection**

## 2.2.3 Habitat Assessment

The vegetation and habitats of the USC campus are well studied and support the breeding, foraging and habitation requirements of a diverse assemblage of native fauna species across habitats including wetlands, modified grasslands and fragmented forest communities which have been retained during the development of the area in recent years.

The unirrigated lawn and slashed, isolated tussocks of *Setaria* grass which constitute the vegetation of the proposed link road, is of limited value to any but a small handful of the common, urban-adapted species. The most notable exception to this is the Eastern Grey kangaroo, which utilises these lawns as evidenced by multiple fresh scats observed there during our site visit.

While the proposed works impact area is providing habitat for Kangaroos onsite it is not considered to be a preferred habitat area for the species within the wider area. USC campus is approximately 100 hectares in site and of this approximately 25-30 hectares may be utilized by the Easter grey kangaroo population for foraging.

The loss of approximately 0.3 ha of kangaroo grazing habitat is not considered a significant impact on the local population as the size of the local kangaroo population is so low that grazing habitat is not a likely limiting resource.

Further, GPS collaring on the USC campus shows that kangaroos prefer to graze on the watered lawns around the centre and north-east of the campus, and slashed *Setaria* paddocks in low lying eastern areas of the campus.

Kangaroos spent relatively little time on dry lawns of the type present in the proposed link road works corridor. This is backed up by the relatively low number (12) of individual kangaroo scats observed throughout the pegged route of the link road, and adjacent lawn. No fauna breeding places were identified in the immediate works area, and given the fact of it being entirely lawn and low slashed grasses, and mostly steep, it is unlikely that any species do use the site for that purpose.

### **2.3 Kangaroo Research at USC**

The campus is home to a much-studied population of *M. giganteus*. This mob of kangaroos are a published case study in the interactions of urban kangaroos with the landscape (Brunton *et al.* 2018a), stress in urban kangaroos (Brunton *et al.* 2019a) and the conservation management of urban kangaroos (Brunton *et al.* 2018b). USC staff and students also regularly undertake surveys of the kangaroo population, and operate an informal network of correspondents who report kangaroo incidents (usually car strikes or dog attacks).

The development of Sippy Downs over the past 11 years, and the increased capacity and use of the road network have devastated the local kangaroo population (Brunton *et al.* 2018a). From 2010 to 2016 the kangaroo population declined from 84 to 16 individuals. The population has hovered around 12 adult kangaroos since then (S. Burnett unpublished data). Human-related deaths, primarily road kill the reason behind this continued low population size.

At the time of writing, the best estimate is that 13 adult grey kangaroos occupy the USC campus, which includes the link road site (Elizabeth Brunton, USC., pers. comm. 6/10/21).

## 3.0 POTENTIAL IMPACTS OF WORKS

### 3.1 *Kangaroo population*

Construction and operation of the link road has the potential to impact Kangaroos and their habitat unless appropriate mitigation strategies are employed. Potential impacts may include the following:

- Injuries to kangaroos when traversing the construction site. At certain phases of construction, kangaroos will potentially be exposed to construction materials and other hazards on which they may be injured. Scats detected during the site visit indicate that kangaroos do currently move through the future construction zone.
- The introduction of another vehicle movement corridor in the form of the link road increases the potential for Kangaroo and vehicle interactions. These can result both in Kangaroo mortality associated with vehicle strike as well as potential for vehicles to have crashes associated with avoidance of Kangaroos. When referring to the Sippy Downs kangaroo population, Brunton *et al.* (2018a) state that the long-term success of urban kangaroo populations is dependent on their ability to maintain home ranges with minimal exposure to traffic. Further they identify that the main risk to kangaroos in the Sippy Downs area is having to cross roads regularly to access food or shelter.
- The positioning of the link road alongside a hard fence (the USC-Chancellor Park boundary) creates a confined space in which startled kangaroos are likely to move erratically, and be prone to leaping in front of moving cars.
- Potential release of sediments and nutrients into adjacent wetland from earthworks if the area is not stabilised or if large rain events occur during works.

Continued modification of Kangaroo habitats within the area combined with the potential for species mortality associated with dog interactions and vehicle strikes remain the primary threats to the continued presence of the species in the area.

### 3.2 *Sustainable Population Limit for USC*

A sustainable population limit (or carrying capacity) is an estimate of the number of kangaroos an area can sustain, without supplementary feeding and watering and while meeting management objectives for the area (DELWP, 2014). The sustainable population limit is expressed as a ratio (kangaroos / ha).

In the Guide to preparing a Kangaroo Management Plan for Melbourne's Growth Corridors, DELWP uses a value of 1.0 kangaroo/ha as a rough guide to estimating the carrying capacity of a Australian grassy woodland site. If this were applied to the approximately 30ha of kangaroo habitat on the USC campus it suggests a carrying capacity of 30 kangaroos, or about twice the current population size.

The management of a population of kangaroos in this area is complex due to the USC campus containing various infrastructure, the increasing urbanisation of Sippy Downs, the presence of arterial roads, and lack of adjacent reserves for mobs to seek refuge. It is noted that the Mooloolah River National Park is across Claymore Road, but that it contains little kangaroo foraging habitat.

## 4.0 KANGAROO MANAGEMENT PLAN

The decline and continued high mortality rate affecting the USC kangaroo mob has led to considerable effort being put into preserving this kangaroo population. Kangaroo passageways linking the campus with Mooloolah River National Park were incorporated into the Claymore Road upgrade in the late 2000s, and the University has actively managed kangaroo threats under its control, and supported research to better understand kangaroo ecology and conservation needs in urbanising landscapes (Brunton *et al.* 2018a, Brunton *et al.* 2018b, Brunton *et al.* 2019a, 2019b, Brunton *et al.* 2020, Hume *et al.* 2019).

While it is not considered that the loss of a small area (approx. 3000m<sup>2</sup>) of sub-optimum Kangaroo grazing area associated with works will impact on the potential survival of the species within the locality it is considered that the potential for further interactions with vehicles is a real and ongoing threat given the low population numbers of Kangaroos present.

It is therefore recommended that appropriate avoidance and mitigation strategies are employed to minimise the potential interaction of vehicles and native fauna, including Kangaroos, as follows:

### 4.1 Design Considerations

1. The alignment of the link road should be placed as close to the existing timber fence as possible in order to reduce the incentive for Kangaroos and other native fauna species to cross the paved road, thereby reducing the potential for vehicle interaction.
2. It is recommended that any areas of land that do remain between the road and the existing fence are not grassed to prevent them providing foraging areas for Kangaroos.
3. Any areas located between the fence could be paved for pedestrian or bicycle use or be mulched and planted with native groundcover species that do not provide foraging opportunities for Kangaroos or create habitat areas for other native fauna.
4. Do not include reticulated irrigation or drainage systems which could improve the quality of Kangaroo forage in proximity to the link road.
5. Do not provide day time shelter for kangaroos. This includes shrubs or trees under which kangaroos can seek shade during the day.

### 4.2 Construction Considerations

1. Prepare a Construction Environmental Management Plan for the proposed works.
2. Works to be undertaken in accordance with Council's Least Concern Fauna Species Management Program.
3. Invite qualified person to present site specific induction for all site personnel which includes the presence of Kangaroo and other native fauna species onsite and the findings of this KMP.



4. The site should have effective temporary construction fencing in place at all stages where fauna injury risks are present. Ensure that gates are closed, and without gaps each evening.
5. No excavated holes should remain open onsite as they have the potential to act as traps for native fauna species and/or may injure kangaroos.
6. If a listed fauna species is identified on site prior to or during works, works to occur in accordance with an approved Species Management Plan for the listed species.

### **4.3 Post-Construction Considerations**

1. Maintain the proposed limited use of the link road i.e. limited speeds and only during school drop off and pickup times.
2. Council to establish signage encouraging the reporting of kangaroo issues on the link road.
3. Council to consider road speed reductions within the area to reduce potential for car strikes.
4. Council to consider fauna signage on local roads to inform residents and road users to drive with care.
5. Council to undertake an education program for local residents, regarding dog management in the area i.e. keeping dogs on leash, ensuring dogs are not roaming at night, microchipping dogs, etc. in relation to protection of native fauna species.
6. Council to review wildlife corridors in the area and determine if more wildlife corridors can be established to facilitate kangaroo movement into and out of the USC site for greater gene flow and population augmentation.
7. Monitoring of the movement and foraging patterns of the remaining Kangaroo population onsite should be continued by USC and SCC in order to understand how the operation of the link road impacts on the species. If it is determined that the operation of the link road is having a negative impact on the population or is directly responsible for Kangaroo mortality, adaptive management strategies to reduce detrimental impacts should be employed.
8. Maintaining attributes of the road and landscaped verges as per **Section 4.2** above to maintain the site to be as unappealing to kangaroos as possible.

The implementation of design and construction strategies to avoid and mitigate potential impacts on Kangaroo populations remaining in the area combined with implementation of the existing management protocols developed by USC may assist in preserving this vulnerable Kangaroo population.

## 5.0 SUMMARY

The assessment of the proposed works corridor for the link road between Scholar Drive and Parkville Road found:

- The proposed works corridor does not require removal of native vegetation.
- The Eastern Grey Kangaroo (*Macropus giganteus*) population in Sippy Downs may use this grassy area for foraging and movement
- Interactions with vehicles within the area are the main threat to the kangaroo population on the USC campus.
- The corridor does not contain significant habitat features for other native fauna.
- The works would result in a loss of approximately 3000m<sup>2</sup> of grassed areas, which due to lack of watering, is sub-optimum grazing for the kangaroo population on site.
- The works would not directly impact on the kangaroo population by removing any limiting resource.
- The works and ongoing operation of the Link Rd have the potential to add another source of mortality to the threatened kangaroo mob which inhabit the USC campus although this should be mitigated by the vehicle movement parameters i.e. reduced speed and limited opening times as well as appropriate design modifications to reduce the occurrence of the species adjacent to vehicle movement corridors.

---

## REFERENCES

Brunton, B., Srivastava, S., and Burnett, S. 2018a. Spatial ecology of an urban eastern grey kangaroo (*Macropus giganteus*) population: local decline driven by kangaroo–vehicle collisions. *Wildlife Research* 45(8) 685-695 <https://doi.org/10.1071/WR18077>

Brunton, B., Srivastava, S., Schoeman, D., Burnett, S. 2018b. Quantifying trends and predictors of decline in eastern grey kangaroo (*Macropus giganteus*) populations in a rapidly urbanising landscape. *Pacific Conservation Biology*. Vol. 24, No. 1, pp.63-73

Brunton, E., Clemente, C and Burnett, S. 2019a. Not all urban landscapes are the same: interactions between urban land use and stress in a large herbivorous mammal. *Ecological Applications* <https://doi.org/10.1002/eap.2055>

Brunton, E., Bolin, J., Leon, J., and Burnett, S. 2019b. Fright or flight? Behavioural responses of kangaroos to drone-based monitoring. *Drones* 3(2), 41-52.

Brunton, E., Leon, J., and Burnett, S. 2020. Evaluating the efficacy and optimal deployment of thermal infrared and true-colour imaging when using drones for wildlife monitoring. *Drones*.

Department of Environment, Land, Water and Planning Melbourne (Victorian Government). 2014. Guide to preparing a Kangaroo Management Plan for Melbourne's Growth Corridors.

Hume, G. Brunton, E., Burnett, S. 2019. Eastern grey kangaroo (*Macropus giganteus*) vigilance varies between human-modified and natural environments. *Animals*. 9(8).

## APPENDIX A – INCIDENT AND COMPLAINT FORM

Incident Reference No.		Nature of Incident	
		<input type="checkbox"/> Fauna <input type="checkbox"/> Habitat <input type="checkbox"/> Water quality <input type="checkbox"/> Other – Specify	
<b>PROJECT SITE</b>			
Project Name:			
Project Manager:			
Project Manager's Contact Details:			
<b>INCIDENT DETAILS</b>			
Submitter's Name:			
Submitter's Contact Details:			
Date of Incident:		Time	
Location of Incident:			
Description of Incident			
Reported to – Name:		Position	
Date Reported:		Time	
<b>CORRECTIVE ACTIONS IMPLEMENTED</b>			
<b>FURTHER ACTION REQUIRED?</b>			
Did the incident have potential for off-site environmental harm?		<input type="checkbox"/> Yes <input type="checkbox"/> No	
Details of potential harm			
Did correction action stop environmental harm?		<input type="checkbox"/> Yes <input type="checkbox"/> No	
Is further corrective action or assessment required?		Attach Details	
Does the incident require referral to regulatory authorities?		<input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Unsure	
If so has referral occurred?		Date	
<b>SIGN OFF</b>			
All actions completed satisfactorily		<input type="checkbox"/> Yes <input type="checkbox"/> No	
Signature		Date	
Name		Position	



## APPENDIX B – Fauna recorded on USC campus

SCIENTIFIC NAME	COMMON NAME
<b>AMPHIBIANS</b>	
<i>Bufo marinus</i> *	Cane Toad
<i>Crinia tinnula</i>	Wallum Froglet
<i>Limnodynastes peroni</i>	Striped Marsh Frog
<i>Litoria caerulea</i>	Common Tree Frog
<i>Litoria fallax</i>	Eastern Sedgefrog
<i>Litoria gracilenta</i>	Graceful Tree Frog
<i>Litoria nasuta</i>	Striped-Rocket Frog
<i>Litoria rubella</i>	Naked Tree Frog
<i>Pseudophryne raveni</i>	Raven's Brood Frog
<b>BIRDS</b>	
<i>Acanthiza pusilla</i>	Brown Thornbill
<i>Accipiter cirrhocephalus</i>	Collared Sparrowhawk
<i>Accipiter fasciatus</i>	Brown Goshawk
<i>Accipiter novaehollandiae</i>	Grey Goshawk
<i>Acridotheres tristis</i> *	Common Myna*
<i>Acrocephalus stentoreus</i>	Clamorous Reed-Warbler
<i>Anas rhyncotis</i>	Australian Shoveler
<i>Anas superciliosa</i>	Pacific Black Duck
<i>Anhinga melanogaster</i>	Darter
<i>Anseranas semipalmata</i>	Maggie Goose
<i>Anthochaera chrysoptera</i>	Little Wattlebird
<i>Anthus novaeseelandiae</i>	Richard'S Pipit
<i>Ardea alba</i>	Great Egret
<i>Ardea ibis</i>	Cattle Egret
<i>Ardea intermedia</i>	Intermediate Egret
<i>Ardea pacifica</i>	White-Necked Heron
<i>Artamus cyanopterus</i>	Dusky Woodswallow
<i>Artamus leucorhynchus</i>	White-Breasted Woodswallow
<i>Aviceda subcristata</i>	Pacific Baza
<i>Aythya australis</i>	Hardhead
<i>Cacatua galerita</i>	Sulphur-Crested Cockatoo
<i>Cacatua roseicapilla</i>	Galah
<i>Cacatua sanguinea</i>	Little Corella
<i>Cacomantis flabelliformis</i>	Fantailed Cuckoo
<i>Calyptorhynchus funereus</i>	Yellow-Tailed Black-Cockatoo
<i>Centropus phasianinus</i>	Pheasant Coucal
<i>Chenonetta jubata</i>	Australian Wood Duck
<i>Chlidonias hybridus</i>	Whiskered Tern

<i>Chrysococcyx lucidus</i>	Shining Bronze Cuckoo
<i>Chrysococcyx minutillus</i>	Little Bronze Cuckoo
<i>Circus approximans</i>	Swamp Harrier
<i>Circus assimilis</i>	Spotted Harrier
<i>Cisticola exilis</i>	Golden-Headed Cisticola
<i>Colluricincla harmonica</i>	Grey Shrike-Thrush
<i>Colluricincla megarhyncha</i>	Little Shrike-Thrush
<i>Coracina novaehollandiae</i>	Black-Faced Cuckoo-Shrike
<i>Coracina tenuirostris</i>	Cicadabird
<i>Corvus orru</i>	Torresian Crow
<i>Coturnix ypsilophora</i>	Brown Quail
<i>Cracticus nigrogularis</i>	Pied Butcherbird
<i>Cracticus torquatus</i>	Grey Butcherbird
<i>Cygnus atratus</i>	Black Swan
<i>Dacelo novaeguineae</i>	Laughing Kookaburra
<i>Dendrocygna arcuata</i>	Wandering Whistling-Duck
<i>Dendrocygna eytoni</i>	Plumed Whistling-Duck
<i>Dicaeum hirundinaceum</i>	Mistletoebird
<i>Dicrurus bracteatus</i>	Spangled Drongo
<i>Egretta garzetta</i>	Little Egret
<i>Egretta novaehollandiae</i>	White-Faced Heron
<i>Elanus axillaris</i>	Black-Shouldered Kite
<i>Elseyornis melanops</i>	Black-Fronted Dotterel
<i>Entomyzon cyanotis</i>	Blue-Faced Honeyeater
<i>Eopsaltria australis</i>	Eastern Yellow Robin
<i>Eudynamys scolopacea</i>	Eastern Koel
<i>Falco berigora</i>	Brown Falcon
<i>Falco cenchroides</i>	Nankeen Kestrel
<i>Falco longipennis</i>	Australian Hobby
<i>Fulica atra</i>	Eurasian Coot
<i>Gallinago hardwickii</i>	Latham's Snipe
<i>Gallinula tenebrosa</i>	Dusky Moorhen
<i>Gallirallus philippensis</i>	Buff-Banded Rail
<i>Geopelia humeralis</i>	Bar-Shouldered Dove
<i>Gerygone olivacea</i>	White-Throated Gerygone
<i>Glossopsitta concinna</i>	Musk Lorikeet
<i>Grallina cyanoleuca</i>	Magpie-Lark
<i>Gymnorhina tibicen</i>	Australian Magpie
<i>Haliaeetus leucogaster</i>	White-Bellied Sea-Eagle
<i>Haliastur indus</i>	Brahminy Kite
<i>Haliastur sphenurus</i>	Whistling Kite
<i>Himantopus himantopus</i>	Black-Winged Stilt
<i>Hirundapus caudacutus</i>	White-Throated Needletail
<i>Hirundo ariel</i>	Fairy Martin

<i>Hirundo neoxena</i>	Welcome Swallow
<i>Hirundo nigricans</i>	Tree Martin
<i>Irediparra gallinacea</i>	Comb-Crested Jacana
<i>Lalage leucomela</i>	Varied Triller
<i>Lichenostomus chrysops</i>	Yellow-Faced Honeyeater
<i>Lichmera indistincta</i>	Brown Honeyeater
<i>Lonchura castaneothorax</i>	Chestnut-Breasted Mannikin
<i>Macropygia amboinensis</i>	Brown Cuckoo Dove
<i>Malurus lamberti</i>	Variegated Fairy-Wren
<i>Malurus melanocephalus</i>	Red-Backed Fairy-Wren
<i>Manorina melanocephala</i>	Noisy Miner
<i>Megalurus timoriensis</i>	Tawny Grassbird
<i>Meliphaga lewinii</i>	Lewin's Honeyeater
<i>Melithreptis albogularis</i>	White-Throated Honeyeater
<i>Merops ornatus</i>	Rainbow Bee-Eater
<i>Microeca fascinans</i>	Jacky Winter
<i>Milvus migrans</i>	Black Kite
<i>Myiagra rubecula</i>	Leaden Flycatcher
<i>Myzomela sanguinolenta</i>	Scarlet Honeyeater
<i>Neochmia temporalis</i>	Red-Browed Finch
<i>Nettapus coromandelianus</i>	Cotton Pygmy-Goose
<i>Nycticorax caledonicus</i>	Nankeen Nightheron
<i>Ocyphaps lophotes</i>	Crested Pigeon
<i>Oriolus sagittatus</i>	Olive-Backed Oriole
<i>Pachycephala pectoralis</i>	Golden Whistler
<i>Pachycephala rufiventris</i>	Rufous Whistler
<i>Pandion haliaetus</i>	Eastern Osprey
<i>Pardalotus striatus</i>	Striated Pardalote
<i>Passer domesticus</i>	House Sparrow
<i>Pelecanus conspicillatus</i>	Australian Pelican
<i>Phalacrocorax melanoleucos</i>	Little Pied Cormorant
<i>Phalacrocorax sulcirostris</i>	Little Black Cormorant
<i>Philemon citreogularis</i>	Little Friarbird
<i>Philemon corniculatus</i>	Noisy Friarbird
<i>Phylidonyris nigra</i>	White-Cheeked Honeyeater
<i>Platalea regia</i>	Royal Spoonbill
<i>Platycercus adscitus</i>	Pale-Headed Rosella
<i>Podargus strigoides</i>	Tawny Frogmouth
<i>Podiceps cristatus</i>	Great Crested Grebe
<i>Porphyrio porphyrio</i>	Purple Swamphen
<i>Porzana pusilla</i>	Bailon's Crake
<i>Psophodes olivaceus</i>	Eastern Whipbird
<i>Rhipidura fuliginosa</i>	Grey Fantail
<i>Rhipidura leucophrys</i>	Willie Wagtail

<i>Sericornis frontalis</i>	White-Browed Scrubwren
<i>Sphecotheres viridis</i>	Figbird
<i>Strepera graculina</i>	Pied Currawong
<i>Streptopelia chinensis*</i>	Spotted Turtle-Dove*
<i>Sturnus vulgaris</i>	Common Starling*
<i>Symposiachrus trivirgatus</i>	Spectacled Monarch
<i>Tachybaptus novaehollandiae</i>	Australasian Grebe
<i>Taeniopygia bichenovii</i>	Double-Barred Finch
<i>Threskiornis molucca</i>	Australian White Ibis
<i>Threskiornis spinicollis</i>	Straw-Necked Ibis
<i>Todiramphus macleayii</i>	Forest Kingfisher
<i>Todiramphus sanctus</i>	Sacred Kingfisher
<i>Trichoglossus chlorolepidotus</i>	Scaly-Breasted Lorikeet
<i>Trichoglossus haematodus</i>	Rainbow Lorikeet
<i>Vanellus miles novaehollandiae</i>	Masked Lapwing
<i>Zosterops lateralis</i>	Silvereye
<b>MAMMALS</b>	
<i>Antechinus flavipes</i>	Yellow-Footed Antechinus
<i>Austronomus australis</i>	White-Striped Mastiff Bat
<i>Canis lupus dingo</i>	Dingo
<i>Chalinolobus gouldii</i>	Gould'S Wattled Bat
<i>Felis catus</i>	Domestic Cat
<i>Hydromys chrysogaster</i>	Water Rat
<i>Isodon macrourus</i>	Northern Brown Bandicoot
<i>Lepus capensis</i>	Brown Hare
<i>Macropus giganteus</i>	Eastern Grey Kangaroo
<i>Melomys burtoni</i>	Grassland Melomys
<i>Melomys cervinipes</i>	Fawn-Footed Melomys
<i>Miniopterus australis</i>	Little Bentwing Bat
<i>Miniopterus oceanensis</i>	Common Bentwing Bat
<i>Mormopterus beccari</i>	Beccari's Freetail Bat
<i>Mormopterus ridei</i>	Eastern Freetail Bat
<i>Mus musculus</i>	House Mouse
<i>Myotis adversus</i>	Fishing Bat
<i>Nyctophilus sp.</i>	Long-Eared Bat Sp.
<i>Petaurus norfolcensis</i>	Squirrel Glider
<i>Planigale maculata</i>	Common Planigale
<i>Pseudocheirus peregrinus</i>	Common Ringtail Possum
<i>Pteropus alecto</i>	Black Fruitbat
<i>Pteropus poliocephalus</i>	Grey-Headed Fruitbat
<i>Rattus lutreolus</i>	Swamp Rat
<i>Rattus rattus</i>	Black Rat
<i>Rattus tunneyi</i>	Pale Field Rat



<i>Scoteanax/Scotorepens orion</i>	Greater Or Eastern Broad-Nosed Bat
<i>Scotorepens sp./Chalinolobus nigrogriseus</i>	Central Easterbroad-Nosed Bat Or Hoary Wattled Bat
<i>Trichosurus vulpecula</i>	Common Brushtailed Possum
<i>Vulpes vulpes*</i>	Red Fox*
<b>REPTILES</b>	
<i>Anomolopus verreauxii</i>	Vereaux's Skink
<i>Cyclodomorphus gerrardii</i>	Pink-Tongued Skink
<i>Demansia psammophis</i>	Yellow-Faced Whip-Snake
<i>Demansia vestigiata</i>	Lesser Black Whip Snake
<i>Eulamprus quoyii</i>	Golden Water Skink
<i>Hemidactylus frenatus</i>	Asian House Gecko
<i>Lampropholis amicula</i>	Secretive Shade Skink
<i>Lampropholis delicata</i>	Garden Skink
<i>Lampropholis guichenoti</i>	Grass Skink
<i>Lialis burtonis</i>	Burton's Legless Lizard
<i>Morelia spilota</i>	Carpet Python
<i>Physignathus leseuerii</i>	Eastern Water Dragon
<i>Pogona barbata</i>	Eastern Bearded Dragon
<i>Pseudechis porphyriaceus</i>	Red-Bellied Black Snake
<i>Pseudonaja textilis</i>	Eastern Brown Snake
<i>Pygopus lepidopodus</i>	Eastern Scaly-Foot
<i>Tiliqua scincoides</i>	Blue-Tongued Lizard
<i>Tropidonophis mairii</i>	Freshwater Snake
<i>Varanus varius</i>	Lace Monitor
<i>Wollumbinia latisternum</i>	Saw-Shelled Turtle

KEY      V- Vulnerable      \* Introduced

## APPENDIX C - RELOCATION PROCEDURE

OBJECTIVE OF RELOCATION PROCEDURE:	TO PREVENT INJURY OR DEATH TO NATIVE FAUNA FOUND WITHIN THE CLEARING/ CONSTRUCTION ZONE PRIOR TO OR DURING WORKS.
<b>Steps to follow in the event of relocation:</b>	
1.	Work will cease in the immediate vicinity of the sighting until the animal has moved off or a suitably qualified Fauna Spotter-catcher has removed it.
2.	All fauna species 'not of concern' will be managed by the Fauna Spotter-catcher and all personnel working on site in accordance the <i>Animal Care and Protection Act 2001</i> , <i>Nature Conservation Act 1992</i> and the <i>Code of Practice: Care of Sick, Injured or Orphaned Protected Animals in Queensland</i> .
3.	If any EVNT listed fauna are found onsite during clearing or construction, work will immediately cease in the vicinity of the sighting, EHP will be notified and a Species Management Plan will be prepared.
4.	If the animal appears to be healthy, a relocation zone within an adjacent area of suitable, undisturbed habitat will be determined by the Ecologist/Fauna Spotter-catcher.
5.	Care will be taken to ensure that aquatic and amphibian specimens are kept moist and free of disease following the procedures contained in the <i>Interim hygiene protocol for handling amphibians</i> (EHP, 2013, <b>Appendix C</b> ). Disposable gloves will be worn when handling frogs. To prevent cross-contamination, new gloves and a clean plastic bag will be used for each frog specimen. Unnecessary handling will be avoided and specimens released as soon as possible, to minimise stress incurred.
6.	In the event a frog appears to be sick, or is dead, the procedures for handling sick or dead frogs outlined in the <i>Interim hygiene protocol for handling amphibians</i> will be followed (EHP, 2013, <b>Appendix C</b> ). Frogs exhibiting one or more of the symptoms for sick frogs and considered unlikely to survive transportation will be euthanased. Typical clinical signs of frogs with Chytridiomycosis include lethargy, inappetence, skin discoloration, presence of excessive sloughed skin, and sitting unprotected during the day with hind legs held loosely to the body. Details of sick or dead listed frogs found on the project site will be recorded and reported to SCC/EHP using the Sick or Dead Frog Collection Form template ( <b>Appendix E</b> ).
7.	Any animal requiring care or treatment will be immediately transported to a veterinarian or licensed wildlife carer.
8.	Specimens will be relocated into appropriate habitat features to prevent predation from other fauna.
9.	Fauna should be released at a time and place that will not endanger it (i.e. release should not increase the risk of stress or predation to the species).
10.	Details of fauna relocations will be recorded using the Translocation Record Form template ( <b>Appendix E</b> ).

Native Foresters