

Koala occupancy surveys in the *Caloundra Transport Corridor Upgrade* area



Prepared for Sunshine Coast Council

by Scientific Koala Surveys / Detection Dogs for Conservation

Dr Romane Cristescu & Kye McDonald

Disclaimer

This report was prepared in accordance with the scope of work agreed with the Sunshine Coast Council and is subject to the specific time, cost and other constraints as defined by the scope of work. To prepare this report, Scientific Koala Surveys and Detection Dogs for Conservation relied on information supplied by Sunshine Coast Council, and does not accept responsibility for the accuracy or completeness of this information. Scientific Koala Surveys / Detection Dogs for Conservation also relied on information gathered at particular times and under particular conditions, and does not accept responsibility for any changes or variances to this information which may have subsequently occurred. Accordingly, the authors of the report provide no guarantee, warranty or representation in respect to the accuracy, adequacy or completeness of the information, whether generally or for use or reliance in specific circumstances. To the extent permitted by law, the authors exclude any liability, including any liability for negligence, for any loss, damage, injury, illness howsoever caused, including (with limitation) by the use of, or reliance upon, the information, and whether arising from errors or omissions or otherwise.

This report is subject to copyright protection and the copyright owner reserves its rights.

Citation

Romane Cristescu & Kye McDonald, Scientific Koala Surveys / Detection Dogs for Conservation. Koala occupancy surveys in the Caloundra Transport Corridor Upgrade area. Prepared for the Sunshine Coast Council. March 2025. 29p.

Acknowledgements

We acknowledge the Jinibara and Kabi Kabi people of this land where we work, play and live, and where this research was conducted. We acknowledge their spiritual and cultural belonging and recognise their continuing connection to land, waters and culture. We pay our respects to Elders past, present and emerging.

Document version control

Version	Date	Authors	Reviewers
V1	25/02/2025	Romane Cristescu	Kye McDonald
V2	16/03/2025	Kye McDonald	Romane Cristescu

Table of Contents

Koala occupancy surveys in the <i>Caloundra Transport Corridor Upgrade</i> area.....	1
Disclaimer.....	2
Citation.....	2
Acknowledgements.....	3
Document version control	3
Table of Contents.....	4
1 Introduction	5
1.1 Background	5
1.2 Scope of works.....	5
2 Methods	6
2.1 Koala occupancy using bioacoustics	6
2.2 Koala occupancy using detection dogs	7
3 Results	9
3.1 Koala occupancy using bioacoustics	9
3.2 Koala occupancy using detection dogs	11
4 Discussion.....	13
References	15
Appendices.....	17
Appendix A - Bioacoustic recorder deployment locations, with recording times and durations.....	17
Appendix B - Pictures of survey sites	18
End of report.....	29

1 Introduction

1.1 Background

The koala (*Phascolarctos cinereus*) is a highly specialised, folivorous, arboreal marsupial, recognised internationally as one of Australia's most iconic species (Martin and Handasyde 1990). Bringing more than three billion dollars to the Australian economy each year (Conrad 2014, Markwell 2021), the koala is a culturally significant species that serves as an ecological umbrella for a wide range of wildlife inhabiting forest ecosystems (Ward, Rhodes et al. 2020). Koalas are facing multiple threats, including habitat loss, degradation and fragmentation, disease, climate change (including worsening bushfires), vehicle strikes and canine attacks (Melzer, Carrick et al. 2000, McAlpine, Rhodes et al. 2006, Gonzalez-Astudillo, Allavena et al. 2017, Beyer, de Villiers et al. 2018, Shabani, Shafapourtehrany et al. 2023). In recent decades, koala populations have experienced dramatic declines which legislation has so far failed to reverse (McAlpine, Lunney et al. 2015, Tisdell, Preece et al. 2017). Following the 2019-2020 Black Summer megafire, the species was listed as Endangered in Queensland, New South Wales and the Australian Capital Territory under the Australian Environment Protection and Biodiversity Conservation (EPBC) Act 1999 (Commonwealth of Australia 2022).

1.2 Scope of works

This project aimed to determine koala occupancy in areas of interest for the Caloundra Transport Corridor Upgrade, within the Sunshine Coast Council (SCC), as identified in:

- Caloundra City Council Report for Koala Habitat Survey and Mapping. Final Report May 2008. GHD
- Caloundra Transport Corridor Upgrade (Figure 7.5: Potential habitat mapping for the Koala). Aurecon
- Sunshine Coast Council Drone Survey, Thermal Drone Koala Survey | Technical Report May 2023. 2023 Halo Nature Reserves Pty Ltd T/A Endeavour Vet Ecology

Two koala survey methods were selected to assess koala occupancy: bioacoustics and koala scat detection dog. Bioacoustics is a powerful way, during the koala breeding season, to detect the presence of male koalas through their periodic bellowing (Law, Gonsalves et al. 2020). Detection dog surveys are more accurate and efficient than human-led surveys to locate koala scats (Cristescu, Foley et al. 2015) and can be adapted to a range of koala survey types, including for occupancy surveys (Cristescu, Miller et al. 2019). For occupancy surveys, the added benefit of scats is that they can remain present and detectable within the survey area for months, therefore enabling surveys to encompass a longer sampling period (compared, for instance, to bioacoustics which sample only the deployment period).

2 Methods

2.1 Koala occupancy using bioacoustics

Data was recorded using Song Meter Micro (SMM, Wildlife Acoustics, Maynard, MA, USA) acoustic recorders at 14 sites within the area of interest (see [Appendix A](#) for coordinates, and [Appendix B](#) for site pictures). Acoustic recorders were programmed to record sounds from sunset until sunrise, the peak time for koala calls (Ellis, Bercovitch et al. 2011). Song Meter Micros have sampling rates of 32,000 Hz, resolution of 16 bits per sample and 24 dB gain and can typically detect koala bellows from a distance of up to 150 metres. These song meters were fitted with a 64-microSD card and Eneloop NiMH batteries, before being attached to a tree trunk at a height of 1.5 m above ground using lockable bands or rope. Typically, these recorders can operate for between 9 and 12 nights on a single deployment before batteries and microSD card require replacing if longer monitoring periods are sought.

Raw audio recordings were scanned with the open-source AviaNZ acoustic software (Marsland, Priyadarshani et al. 2019) using a koala recogniser (version: DPI_Male_Koala_V3_CNN15_10-02-23, Department of Primary Industries, Towsey, Planitz et al. 2012) developed using a deep learning procedure (Himawan, Towsey et al. 2018). Recordings matched by the recogniser with koala bellows were then validated manually by

visualisation of spectrograms of the audio in AviaNZ (**Figure 1:**), in conjunction with playback of recordings, to delete any false positives.




Figure 1: Examples of spectrograms from AviaNZ; left two images showing a male koala bellow, image on the right showing a false positive (manually marked with a cross by a human verifier).

2.2 Koala occupancy using detection dogs

The detection dog used for these surveys, Austin (**Figure 1:**), is a border-collie cross, trained and tested by the DDC to detect koala scats of any age. Austin is highly experienced in detecting koala scats and has conducted more than 300 surveys.

Upon arrival at each survey site, ecological characteristics that might have influenced the detectability and decay of scats were recorded as rapid scat decay type e.g. wet areas or recent fire (Cristescu, Goethals et al. 2012). The detection dog was fitted with a GPS collar that recorded time and location data, and was directed to search for koala scats. The team also visually searched for any signs of koalas.



Figure 2: Austin, koala scat detection dog.

Handlers are required to visually confirm scat identification and age – for this reason, it is critical that an ecologist or experienced koala researcher is involved. In fact, scat identification presents challenges even to experienced ecologists (Harrington, Harrington et al. 2010). Typical koala scats (**Figure 3**) have the following characteristics (Triggs 1996):

- symmetrical and bullet-shaped (not jelly-bean shaped),
- generally, about 1.5 cm long by 0.5 cm wide (adult koala scat size),
- evenly-sized, fine particles, and
- absence of insect parts (koalas do not eat insects).



Figure 3. Typical characteristics of koala scats.

3 Results

3.1 Koala occupancy using bioacoustics

Acoustic data was recorded over the period 6 February 2025 to 18 February 2025. All 14 deployed acoustic recorders successfully recorded at least 11 full nights of data each. Screening of these recordings resulted in no koala bellow detections at any of the 14 sites during the deployment period (**Figure 4, Appendix A**).




Figure 4: Acoustic recorder deployment locations (red points), all without any koala bellow detections during the monitoring period.

3.2 Koala occupancy using detection dogs

Detection dog surveys occurred on the 7th, 10th, 11th and 18th February 2025, totalling 30.4 km of searched transects, and resulted in no koala scats being detected by the detection dog (**Figure 4**). Some areas could not be covered due to human habitations or being flooded (where scats decay rapidly and cannot efficiently be searched).



Figure 5: Detection dog survey search tracks, with no koala scats detected.

4 Discussion

Appropriate koala survey effort and coverage (**Figure 4-5**) has found no evidence of recent use by koalas within the study area. This is consistent with previous thermal imaging drone surveys which also found no koalas at the time of survey (EVE, 2023), but our results further increase confidence, both by expanding the search area and extending the period of sampling (bioacoustics 11 nights, and detection dog up to a few months).

Like any ecological survey, failure to detect koalas in an area is not necessarily conclusive, and may suggest, for a koala scat survey, either of the following:

- Koalas are not present in the area (i.e., true absence) at the time of the survey. Note that true current absence does not infer that the site has not been used in the past, or could not be used in the future (i.e., it could still be potential koala habitat).
- Koalas occur in the area; however, scats were not detected (i.e., false negative) because:
 - Scats were present at some stage but decayed and disappeared from the environment before the survey was conducted.
 - The dog did not detect the scat; and/or,
 - The dog indicated the presence of a scat, but it was too decayed to be confirmed.

Detection dog surveys can also be hampered by wet conditions, which degrade koala scats faster than normal. In addition, impenetrable and/or flooded bushland for detection dog teams, reduces accessibility and therefore the effectiveness of surveys. Due to the location, multiple sites had water inundation/swampy conditions. The relevant dog track maps in **Figure 4** reflect this, with certain areas within the designated sites not completely covered by the dog team.

However, in combination with the bioacoustics monitoring, complete koala survey coverage was achieved across the entire study area, with no evidence of koala occupation found. It is important to note that, although these survey methods have the power to confidently



Detection Dogs
for Conservation

Scientific Koala Surveys

ascertain presence, confirming absence is difficult. To infer true absence, multiple surveys are generally necessary (MacKenzie and Royle 2005).

References

- Beyer, H. L., D. de Villiers, J. Loader, A. Robbins, M. Stigner, N. Forbes and J. Hanger (2018). "Management of multiple threats achieves meaningful koala conservation outcomes." Journal of Applied Ecology **55**(4): 1966-1975.
- Commonwealth of Australia (2022). "List of Threatened Species Amendment (*Phascolarctos cinereus* (combined populations of Queensland, New South Wales and the Australian Capital Territory)) Instrument 2022. F2022L00131.
- Conrad, E. (2014). "The economic value of the koala." www.savethekoala.com.
- Cristescu, R., K. Goethals, P. B. Banks, F. Carrick and C. Frère (2012). "Persistence and detectability of fecal pellets in different environment and the implication for pellet based census of fauna." International Journal of Zoology 2012, Article ID 631856: doi:10.1155/2012/631856.
- Cristescu, R. H., E. Foley, A. Markula, G. Jackson, D. Jones and C. Frère (2015). "Accuracy and efficiency of detection dogs: a powerful new tool for koala conservation and management." Scientific Reports **5**(1): 8349.
- Cristescu, R. H., R. L. Miller, A. J. Schultz, L. Hulse, D. Jaccoud, S. Johnston, J. Hanger, R. Booth and C. H. Frère (2019). "Developing noninvasive methodologies to assess koala population health through detecting *Chlamydia* from scats." Molecular Ecology Resources **19**(4): 957-969.
- Department of Primary Industries. "Fauna identification service." Retrieved 15/12/2022, 2022, from <https://www.dpi.nsw.gov.au/forestry/science/forest-ecology/fauna-identification-service>.
- Ellis, W., F. Bercovitch, S. FitzGibbon, P. Roe, J. Wimmer, A. Melzer and R. Wilson (2011). "Koala bellows and their association with the spatial dynamics of free-ranging koalas." Behavioral Ecology **22**(2): 372-377.
- Gonzalez-Astudillo, V., R. Allavena, A. McKinnon, R. Larkin and J. Henning (2017). "Decline causes of Koalas in South East Queensland, Australia: a 17-year retrospective study of mortality and morbidity." Scientific reports **7**(1): 1-11.
- Harrington, L. A., A. L. Harrington, J. Hughes, D. Stirling and D. W. Macdonald (2010). "The accuracy of scat identification in distribution surveys: American mink, *Neovison vison*, in the northern highlands of Scotland." European Journal of Wildlife Research **56**: 377-384.
- Himawan, I., M. Towsey, B. Law and P. Roe (2018). "Deep learning techniques for koala activity detection." Proceedings of the Annual Conference of the International Speech Communication Association, Interspeech.
- Law, B., L. Gonsalves, R. Bilney, J. Peterie, R. Pietsch, P. Roe and A. Truskinger (2020). "Using passive acoustic recording and automated call identification to survey koalas in the southern forests of New South Wales." Australian Zoologist **40**(3): 477-486.
- MacKenzie, D. I. and J. A. Royle (2005). "Designing occupancy studies: general advice and allocating survey effort." Journal of applied Ecology **42**(6): 1105-1114.
- Markwell, K. (2021). "Getting close to a national icon: an examination of the involvement of the koala (*Phascolarctos cinereus*) in Australian tourism." Tourism Recreation Research **46**(4): 473-486.

- Marsland, S., N. Priyadarshani, J. Juodakis and I. Castro (2019). "AviaNZ: A future-proofed program for annotation and recognition of animal sounds in long-time field recordings." Methods in Ecology and Evolution **10**(8): 1189-1195.
- Martin, R. and K. Handasyde (1990). "Population dynamics of the koala (*Phascolarctos cinereus*) in southeastern Australia." Biology of the Koala: 75-84.
- McAlpine, C., D. Lunney, A. Melzer, P. Menkhorst, S. Phillips, D. Phalen, W. Ellis, W. Foley, G. Baxter and D. De Villiers (2015). "Conserving koalas: A review of the contrasting regional trends, outlooks and policy challenges." Biological Conservation **192**: 226-236.
- McAlpine, C. A., J. R. Rhodes, J. G. Callaghan, M. E. Bowen, D. Lunney, D. L. Mitchell, D. V. Pullar and H. P. Possingham (2006). "The importance of forest area and configuration relative to local habitat factors for conserving forest mammals: a case study of koalas in Queensland, Australia." Biological Conservation **132**(2): 153-165.
- Melzer, A., F. Carrick, P. Menkhorst, D. Lunney and B. S. T. John (2000). "Overview, Critical Assessment, and Conservation Implications of Koala Distribution and Abundance." Conservation Biology **14**(3): 619-628.
- Shabani, F., M. Shafapourtehrany, M. Ahmadi, B. Kalantar, H. Özener, K. Clancy, A. Esmaili, R. S. da Silva, L. J. Beaumont and J. Llewelyn (2023). "Habitat in flames: How climate change will affect fire risk across koala forests." Environmental Technology & Innovation **32**: 103331.
- Tisdell, C. A., H. J. Preece, S. Abdullah and H. L. Beyer (2017). "Strategies to conserve the koala: cost-effectiveness considerations." Australasian Journal of Environmental Management **24**(3): 302-318.
- Towsey, M., B. Planitz, A. Nantes, J. Wimmer and P. Roe (2012). "A toolbox for animal call recognition." Bioacoustics **21**(2): 107-125.
- Triggs, B. (1996). "Tracks, scats and other traces: A field guide to Australian mammals." Oxford University Press, South Melbourne.
- Ward, M., J. R. Rhodes, J. E. M. Watson, J. Lefevre, S. Atkinson and H. P. Possingham (2020). "Use of surrogate species to cost-effectively prioritize conservation actions." Conservation Biology **34**(3): 600-610.

Appendices

Appendix A - Bioacoustic recorder deployment locations, with recording times and durations

Survey Code	Recorder Name	Latitude	Longitude	First recording	Last recording	Recording duration	Koala bellow detected
AR250206KM	DDC 009	-26.797979	153.103180	06/02/2025 18:38	18/02/2025 03:29	11 days	No
AR250206KM	DDC 010	-26.789486	153.114304	06/02/2025 18:38	18/02/2025 04:30	11 days	No
AR250206KM	DDC 011	-26.791712	153.120316	06/02/2025 18:38	18/02/2025 23:28	12 days	No
AR250206KM	DDC 012	-26.792389	153.116013	06/02/2025 18:38	18/02/2025 21:29	12 days	No
AR250206KM	DDC 013	-26.796251	153.115250	06/02/2025 18:38	18/02/2025 22:28	12 days	No
AR250206KM	DDC 014	-26.789246	153.116913	06/02/2025 18:38	17/02/2025 22:29	11 days	No
AR250206KM	DDC 022	-26.800808	153.113556	06/02/2025 18:38	18/02/2025 05:29	11 days	No
AR250206KM	DDC 023	-26.808767	153.116150	06/02/2025 18:38	18/02/2025 19:28	12 days	No
AR250206KM	DDC 024	-26.792270	153.122711	06/02/2025 18:38	18/02/2025 19:28	12 days	No
AR250206KM	DDC 025	-26.794899	153.123764	06/02/2025 18:38	18/02/2025 20:28	12 days	No
AR250206KM	DDC 026	-26.797607	153.124008	06/02/2025 18:38	18/02/2025 03:30	11 days	No
AR250206KM	DDC 027	-26.795822	153.121231	06/02/2025 18:38	18/02/2025 01:29	11 days	No
AR250206KM	DDC 028	-26.801083	153.125229	06/02/2025 18:28	17/02/2025 19:29	11 days	No
AR250206KM	DDC 029	-26.802683	153.127304	06/02/2025 18:28	18/02/2025 01:29	11 days	No

Appendix B - Pictures of survey sites

Town Reserve (1)



Town Reserve (2)





Town Reserve (3)



Town Reserve (4)



Russell Barker Memorial Park





Queen Street Bushland Park



Ben Bennett Bushland Park (1)



Ben Bennett Bushland Park (2)



Ben Bennett Bushland Park (3)



Gosling Park



Bicentennial Park



End of report